

2008, Vol. 2.0




ChemGenes CORPORATION

Experience Nucleic Acid Expertise

New Featured Products

Sulfurizing Reagent
Reverse RNA Synthesis
TOM phosphoramidites
3'-tBDSilyl amidites
L-RNA amidites
7-Deaza ribo Products
8-Methyl ribo G
Universal Support



ChemGenes has been in business for over 25 years and has recently moved into a state of the art facility in Wilmington, MA. ChemGenes has a full scale modernized lab with the facilities to manufacture in bulk while maintaining its high quality. We have added many new products to our original line to facilitate research in the area of biotechnology.

As the market for oligonucleotides continues to grow, ChemGenes remains committed to introducing novel products, while maintaining its existing product mix. We also have the capacity to custom synthesize products on request.

Our quality is guaranteed! We want to assure you that every product is of the highest purity and conforms to the technical data sheet that accompanies it when shipped.

- ChemGenes takes pride in a long history of customer satisfaction in supplying phosphoramidites that have a purity of 98% or better for most phosphoramidites.
- Each lot of Phosphoramidite must pass an established testing criteria before it can be shipped to customers.



Required QC Tests for Most Phosphoramidites

Solubility test

- Amidites completely dissolve in Acetonitrile to make a 0.1M Solution (water<0.004-0.005gm/100ml). Leave no visible particulate matter.

Coupling Efficiency

- The coupling efficiency of ChemGenes phosphoramidite products are 98% or better.

HPLC

- Greater than 98.5% purity by HPLC.

³¹P NMR

- Doublet peak or single peak.
- Position of each peak is known for each phosphoramidite.
- The value between the peaks is calculated and recorded.

UV – The UV test provides 4 values of data:

- The ratio between 250/260 nm.
- The ratio between 260/280 nm.
- Emax position.
- Extinction Coefficient.

MASS Spectrum

- Performed on each product in +ve and -ve mode.


¹H NMR

- Proton NMR analysis is performed for each product.

TLC

- Single or double spot with no other visible impurity on spotting, 0.2mg/spot.
- Single spot or double spot depends on the phosphoramidite.

Sulfurizing Reagent



((Dimethylamino-methylidene)amino)-3H-1,2,4-dithiazoline-3-thione (DDTT) is a new efficient sulfurizing reagent (see Figure 1.) that shows similar performance in oligonucleotide synthesis compared to Beaucage reagent and has greater stability in the formulated solution.




Figure 1.

A number of **1,2,4-dithiazolines** have been described in literature¹⁻² and demonstrated high sulfurizing efficiency in oligonucleotide synthesis. Our data have also shown superior sulfurization by DDTT reagent in solution phase. Unlike Beaucage reagent DDTT did not yield any detectable level of phosphodiester type of by-product.

((Dimethylamino-methylidene)amino)-3H-1,2,4-dithiazoline-3-thione is readily available from ChemGenes Corp. in bulk and formulated solution.

Our recommended formulations are summarized in the table below (Table 1).

Table 1.

Recommended formulations for the synthesis of phosphorothioate containing oligonucleotides and sulfurizing in solution phase.

DDTT Concentration	Solvent formulation v/v	Oxidation time	Comments
0.05 M	Py/ACN – 3:7 Py/THF – 2:8	2.5 min	Formulation used in oligonucleotide synthesis
0.1 M	Py Py/THF – 4:6	2 min	Formulation used in oligonucleotide synthesis
0.05 M	ACN	10 min	Formulation used in solution phase reaction

References:


1. Xu, Q., Musier-Forsyth, K., Hammer R. P. and Barany G. *Nucleic Acid Research*, 1996, 24, 1602-1607.
2. Xu, Q., Barany G., Hammer R. P. and Musier-Forsyth, K. *Nucleic Acid Research*, 1996, 24, 3643-3645.

RNA Synthesis – Reverse Direction



Phosphoramidites for Reverse RNA Synthesis: Applications:

The novel phosphoramidites, **A-n-bz**, **C-n-bz**, **C-n-ac**, **G-n-ac** and **U** are produced with an HPLC purity of greater than 98% and 31P NMR purity greater than 99%. A novel process of reverse **5'→3'** directed synthesis of RNA oligomers has been developed. Using that method we demonstrated high quality RNA synthesis with **coupling efficiency approaching 99%**.



Synthesis of varieties of therapeutic grade RNA and siRNA etc. requires a modification or labeling of 3'-end of an oligonucleotide. The synthesis of 3'-end modified RNA requiring lipophilic, long chain ligands or chromophores, using **3'→5'** synthesis methodology is challenging, requires corresponding solid support and generally results in low coupling efficiency and lower purity of the final oligonucleotide in general because of large amount of truncated sequences containing desired hydrophobic modification. We have approached this problem by developing reverse RNA monomer phosphoramidites for RNA synthesis in **5'→3'**- direction. They lead to very clean oligonucleotide synthesis allowing for introduction of various modifications at the 3'- end cleanly and efficiently.

A vast number of applications are possible for easy attachment at 3'- End of an oligonucleotide.

1. For attachment of bulky molecules at the 3'- end of the RNA, such as cholesterol, long chain aliphatic chains such as C-18, triethylene glycols, hexaethylene glycols. Direct coupling with these amidites can be achieved easily.
2. Attachment of Polyethylene Glycols such as PEG 2000 amidite and PEG 4000 amidites at the 3'- end of the RNA molecule.
3. For easy attachment of 3'- thiol modification. 3'- Disulfides from readily available amidites, via C-3 disulfide, C-6 disulfide.
4. 3'- Biotin attachment via biotin amidite in a single step and avoiding biotin CPG for this purpose.
5. Modification of 3'- end of the sense strand of siRNA. The modification of the overhang of the sense strand (3'- End) of siRNA is not expected to affect targeted mRNA recognition, as the antisense siRNA strand guides target recognition. Useful modification for improvement of delivery of siRNA can be easily designed.

High Purity Synthetic RNA

RNA Synthesis Approach in Reverse Direction. **Leads to Smooth 3'-conjugation of macromolecules to Synthetic RNA. Reverse RNA Synthesis Results in Complete Absence of M+1 Species.**




Figure 1.

Capillary Gel Electrophoresis (CE) of 21- Mer RNA with 3'- Cholesterol-TEG linker. Reverse direction (**5'→3'**) synthesis and HPLC purification. 1 umole scale. Purity; 99.9%.




Figure 2b.

ESI Mass Spectral analysis of 21- mer RNA with 3'- PEG-2000 attachment, purified RNA as shown in Fig. 2a. The synthesis was carried out in reverse direction (**5'→3'** direction). The PEG-2000 was attached as last step via the corresponding phosphoramidite, ChemGenes catalog; CLP-3119.

Calculated Molecular Weight: 8684.1

Observed Molecular Weight: 8681.1

Note: There is a distribution of at least 14 PEG species of the RNA on both sides of the Calculated molecular weight with PEG-2000. Thus species from 8417.1 to 8945.3 are present with a molecular weight difference of a glycol unit (+/- 44).




Figure 2a.

Capillary Gel Electrophoresis (CE) of the HPLC purified 21- mer RNA with 3'- PEG(Poly ethyleneglycol; MW; 2000). Made by Reverse RNA synthesis method (**5'→3'**-direction). Expedite model 8909- 1 umole scale. Purity; 100%


Salient Features:

A vast number of applications are possible for easy attachment at 3'- End of an oligonucleotide.

1. The crude RNA's have much closer impurities (M-1) in the conventional method (**3'→5'**-direction), as compared to reverse RNA synthesis (**5'→3'** direction). Therefore after purification RNA synthesized by reverse RNA synthesis are purer.
2. The feature mentioned above is much more visible in the synthesis of cholesterol attached to 3'- end of RNA. Therefore it is easier to purify RNA with cholesterol at 3'- end synthesized by reverse RNA synthesis.
3. M+1 impurities are essentially absent in the RNA's synthesized by reverse RNA synthesis method. It is postulated that in the molecule, ribonucleoside- 3'- DMT-2'- tBDSilyl-5'- phosphoramidites, the 3'- DMT is not cleaved by 5-ethylthiotetrazole during the coupling time of oligonucleotide chain extension.
4. RNA containing macromolecules at the 3'- end which are generally inaccessible by conventional methods (**3'→5'**) are easily synthesized by reverse RNA synthesis (**5'→3'** direction). These RNA's can be produced in high purity.
5. 3'- PEG RNA (21- mer) was synthesized, and after purification was essentially 100% pure (see figs. 15 & 16).

TOM Phosphoramidites

3'-tBDSilyl RNA Phosphoramidites



Key Advantages:

- **Superior Quality RNA** using 2'-O-TOM protection: No possibility of 2'-5'-linkage.
- Perfected manufacturing process: **prices comparable to TBDMS.**
- **Higher Coupling Efficiency** due to lower steric hindrance: Reduced Coupling Time (2-4 minutes).
- **No base modification or M + 30 observed** (through extensive chemical ionization mass analysis).
- **High Quality Long Chain Oligos.**

Quality Guaranteed:



- Purity greater than 97% by HPLC.
- 31 P NMR purity ranges from 98 -100%.
- UV Spectral data to conform to highest.
- 1 H NMR data to conform.
- Coupling efficiency greater than 98% per step.
- Ideal for Long Chain Oligos.
- TOM Amidites Produced under GMP guidelines.

- Allows the synthesis of 2'-5'-linked oligos.

- RNA 2',5'-duplexes are not substrates of the enzyme RNase. However, they can inhibit the RNaseH mediated cleavage of a natural DNA: RNA substrate.

Useful Applications

- Determine their exact biological role.
- Extend their biological half life.
- Alter the biological activity of the core structure.




Supports			
B	Protection	Catalog #	Pore Size
A	n-acetyl	N-32001-05	500A
		N-32001-10	1000A
C	n-acetyl	N-32002-05	500A
		N-32002-10	1000A
G	n-acetyl	N-32003-05	500A
		N-32002-10	1000A
U	n/a	N-32005-05	500A
		N-32005-10	1000A

Also Available:

Low to high loading CPG supports with TOM-monomer for uniform deprotection of RNA's.


Now Available Bulk Quantities Manufactured Under GMP Guidelines

Licensed from QIAGEN Inc. for therapeutic RNA market.




Adenosine (n-bz) 3'-tBDSilyl
CED OP

Catalog #
ANP-5681




Cytidine (n-bz) 3'-tBDSilyl
CED OP

Catalog #
ANP-5682





Guanosine (n-ibu) 3'-tBDSilyl
CED OP

Catalog #
ANP-5683



Uridine 3'-tBDSilyl
CED OP

Catalog #
ANP-5684




L Deoxy Nucleoside Phosphoramidites & L Ribonucleoside Phosphoramidites (for the synthesis of synthetic L-DNA & L-RNA)


Key Features:

- A synthetic 56-nucleotide b-L-DNA binds with Vasopressin & is an antagonist to Vasopressin (K.P. Williams, X.H. Liu, T.N.M. Schumacher, H.Y. Liu, D.A. Ausiello, P.S. Kim, & D.P. Bartel, PNAS, USA, 94, 11285-11290, 1997).
- L-DNA are useful for studying DNA interactions. The L-DNA are resistant to cleavage by various endo and exonucleases. Furthermore, they are not cleaved by many DNA cleaving drugs. Thus, they can be used for determining specific interactions from non-specific interactions (Urata H, Ueda Y, Usami Y, Akagi M, Journal of the American Chemical Society, 115 (16): 7135-7138 AUG 11 1993).
- L-RNA shows extraordinary stability in intracellular environments. Selective binding of long chain L-RNA to specific biological macromolecules has been searched via combinatorial libraries (Klussmann S, Nolte A, Bald R, Erdmann VA, Furste JP, Nature Biotechnology, 14 (9): 1112-1115 SEP 1996).

L-deoxy nucleoside phosphoramidites



L-2' -tBDSilyl ribo nucleoside phosphoramidites




7- Deaza- Purine-Phosphoramidites:

7- Deaza-modification finds extensive application in molecular biology & design of oligos with 7- deaza-substitution, in place of multiple G's in the sequence.

Key Features:

- To avoid extensive secondary structure formation in oligos and thereby improve targeted hybridization more effectively.
- Antiparallel triple helix formation with double stranded DNA is favored with this modification.


7- Deaza-2'-deoxy A & G Phosphoramidites



7-Deaza deoxy
Adenosine CED OP
Catalog #: **ANP-4815**

7-Deaza deoxy Guanosine
CED OP
Catalog #: **ANP-4857**

7- Deaza-ribo A & G Phosphoramidites:




7-Deaza Adenosine
CED OP
Catalog #: **ANP-7101**

7-Deaza Guanosine
CED OP
Catalog #: **ANP-7301**


7- Deaza-7-iodo-dA and dG Nucleosides:

Key Modifications:

- For conversion of 7-deaza-7-iodo to C-7-Modified-7-deaza-dA and dG-Modified nucleoside Triphosphates
- Extensive application in molecular biology for diagnostics and sequencing.
- 7- position modifications do not interfere in either PCR or oligo hybridizations.
- The 7- deaza-nucleoside phosphates and triphosphates are currently used in DNA sequencing.



7-Deaza-7-iodo deoxy
Adenosine
Catalog #: **DN-2561**



7-Deaza-7-iodo deoxy
Guanosine
Catalog #: **DN-2563**


8-Methyl Guanosine Phosphoramidites

N-Alkylated Phosphoramidites



8-Methyl-2'-dGuanosine & 8-Methyl-rGuanosine:


- Powerful Z-DNA stabilizer
- Reading effects B & Z transformation in oligonucleotides
- Can be incorporated into appropriate positions of synthetic DNA



8-Methyl deoxy Guanosine
CED OP
Catalog #:
ANP-9274

Applications:

- For study of functional role in Gene Expression, transcription control etc.
- Selectivity of Z DNA in protein interactions
- DNA supercoiling modulation
- Selective Targeting of proteins or enzymes
- Aptamer Design and Therapeutic development




8-Methyl ribo Guanosine
CED OP
Catalog #:
ANP-6274


- Due to mutagenic effects of carcinogens, DNA in living organisms is vulnerable to alkylation.
- ChemGenes offers the phosphoramidites for studies of reversal of methylated lesions by use of oligonucleotides incorporating alkylated purine/pyrimidine.

- It has been shown that there is a direct reversal of n-alkylation of methylated bases in oligonucleotides.
- The discovery of an enzyme which is substrate for DNA repair has great implications for repair of such carcinogenic and mutagenic effects. (Ref. 1)


Our featured products include:




N¹-Methyl deoxy
Adenosine
Phosphoramidite
Catalog # **ANP-6121**



N¹-Methyl deoxy
Guanosine
Phosphoramidite
Catalog # **ANP-6122**



N³-Methyl deoxy
Cytidine
Phosphoramidite
Catalog # **ANP-3851**



N³-Methyl Thymidine
Phosphoramidite
Catalog # **ANP-6153**

Ref. 1: (S.C. Trewick, T.F. Henshaw, R.P. Hausinger, T. Lindahl and B. Sedgwick, *Nature*, 419, 174-177, 2002; and another report confirming these observations, P.Falnes, R.F. Johansen, E. Seeberg, *Nature* 419, 178, 2002).

Our Products

Oligo Synthesis Reagents

Natural DNA Amidites & Supports

Ancillary Reagents

Modified DNA Amidites & Supports

Natural RNA Amidites & Supports

Amidites and Supports for Introducing Chromophores & Ligands

Amidites and Supports for 2'-O-Methyl Oligonucleotides

Drying Traps

Oligonucleotide Purification

Nucleosides, Sugars, Purines, & NHS Esters

Unprotected mononucleosides

N-protected mononucleosides

DMT-protected mononucleosides

Phosphoramidite Chemistry Reagents

Sugars & Purines

NHS-Esters

Trisphosphates

Modified Triphosphates

Custom Synthesis

New Featured Products

Universal Support


TOM Amidites

8-Methyl ribo Guanosine Amidite

8-Methyl deoxy Guanosine Amidite

Reverse RNA Synthesis

5'-O-Methyl DNA Amidite



 **ChemGenes**
CORPORATION

33 Industrial Way
Wilmington, MA 01887 USA

Tel: 978-694-4500
Fax: 978-694-4502
Toll Free: 800-762-9323
Email: info@chemgenes.com

www.chemgenes.com